MWARNING FLAMMABLE



Keep away from fire or flame.

To reduce risk of fire, PREP CAREFULLY:

- do not use 26-ml applicator for head and neck surgery or on an area smaller than **8.4 in.** x **8.4 in.** Use a smaller applicator instead.
- solution contains alcohol and gives off **flammable vapors** avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable**.
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair) um of 3 minutes
- do not allow solution to pool
- remove wet materials from prep area

NDC 54365-400-37 NDC 54365-400-38

BD ChloraPrep™ Clear

BD ChloraPrep™ Hi-Lite Orange™

BD ChloraPrep™ Scrub Teal™

26 mL Applicator

2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA) Patient Preoperative Skin Preparation

Not made with natural rubber latex Do Not Reuse

Sterile Solution Applicator is sterile if package is intact

Drug Facts

Active ingredients

Chlorhexidine gluconate 2%

Isopropyl alcohol 70% v/v

Antiseptio Use for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.

Warnings

For external use only. Flammable, keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY:

- do not use 26-ml applicator for head and neck surgery
 do not use on an area smaller than 8.4 in. x 8.4 in. Use a smaller applicator instead
- solution contains alcohol and gives off flammable vapors
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. Wet hair is flammable
- · do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet materials from prep area

Allergy alert:

This product may cause a severe allergic reaction. Symptoms may include:

• wheezing/difficulty breathing

• shock

• facial swelling

• hives

•

If an allergic reaction occurs, stop use and seek medical help right away

③ BD ChloraPrep™ Clear



2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)

Drug Facts (continued)

Do not use

- on patients allergic to chlorhexidine gluconate or any other ingredient in this product
- for lumbar puncture or in contact with the meninges on open skin wounds or as a general skin cleanser

When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- use in a well ventilated area
- maximal treatment area for one applicator is approximately 13.2 in. x 13.2 in. (1126 cm²)
- remove applicator from package; do not touch sponge
 hold the applicator with the sponge down. Pinch wings only once to activate the ampule and release the antiseptic.
- wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skir
- completely wet the treatment area with antiseptic
 dry surgical sites (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds
- moist surgical sites (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes
- do not allow solution to pool; tuck prep towels to absorb solution, and then remove
 allow the solution to completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away.
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available

Other information • store between 15-30 °C (59-86 °F) • avoid freezing and excessive heat above 40 °C (104 °F) • the tint will slowly fade from the skin. Soap and water, or alcohol may be used to remove the tint if desired

Inactive ingredients FD&C green #3 dye (Scrub-Teal applicators only), FD&C yellow #6 dye (Hi-Lite Orange applicators only), USP purified water

Questions? • www.chloraprep.com • call **1-800-523-0502** (M-F 8 a.m.–5 p.m. CST)

CareFusion 213, LLC, El Paso, TX 79912, subsidiary of Becton, Dickinson and Co. • Artwork No. 6-930800 • A050619 • Made in the USA of US and imported parts • www.bd.com

External Use Only Professional Use Only







Purposes

. . Antiseptic

