

Drug Facts (continued)	
Do not use on patients allergic to chlorhexidine gluconate or any other ingredient in this product for lumbar puncture or in contact with the meninges on open skin wounds or as a general skin cleanser 	
When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to en remain. If contact occurs, rinse with cold water right away and contact a doctor.	iter and
Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condit Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.	tion.
Directions • use in a well ventilated area • use in a well ventilated area • maximal treatment area for one applicator is approximately 13.2 in. x 13.2 in. (1126 cm ³) • remove applicator from package; do not touch sponge • hold the applicator with the sponge down. Pinch wings only once to activate the ampule and release the antiseptic. • wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin c completely wet the treatment area with antiseptic • dry surgical sites (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 30 seconds • moist surgical sites (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes • do not allow solution to pool; tuck prep towels to absorb solution, and then remove • allow the solution to completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or • discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It necessary to use the entire amount available.	r wipe away.
Other information • store between 15–30 °C (59–86 °F) • avoid freezing and excessive heat above 40 °C (104 ° • the tint will slowly fade from the skin. Soap and water, or alcohol may be used to remove the tint if desired.	Ϋ́F)