



Prevantics®

Prevent the spread of HAIs with the first and **ONLY 3.15% CHG/70% IPA** formulation with up to 7 days of continued antimicrobial activity.*

*Swabstick and Maxi Swabstick only


 **PDI**®
BE THE DIFFERENCE™

THE CHALLENGE

There were an estimated 722,000 HAIs in U.S. hospitals in 2011, resulting in an estimated 75,000 patient deaths.¹ HAIs affect patients, their families and the people who care for them. They also have an impact on hospital costs and quality. The CDC estimated that CLABSI cost facilities \$16,550 per infection.²

THE SOLUTION

Prevantics® products, designed with 3.15% Chlorhexidine Gluconate (w/v) and 70% Isopropyl Alcohol (v/v) formulation, help prevent the spread of infections and help improve patient outcomes.



Designed for Convenience

Prevantics® products are a convenient delivery system that has no sharp glass to break or activation required.

Prevantics® Applicators

- No glass to break
- Pre-activated
- No waiting
- Easy to open, easy to use
- Pre-moistened
- Swab: Intuitive prep-pad design
- Swabsticks: Unique 2-sided applicator for interdigital areas and skin folds



Prevantics® 1.0 mL Swab



Prevantics® 1.6 mL Swabstick



Prevantics® 5.1 mL Maxi Swabstick

¹CDC HAI Prevalence Survey. Magill SS, Edwards JR, Bamberg W, et al. Multistate Point-Prevalence Survey of Health Care–Associated Infections. N Engl J Med 2014;370:1198-208.

²CDC Vital Signs: Central Line-Associated Blood Stream Infections - United States, 2001, 2008 and 2009. MMWR Weekly, March 4, 2011, 60(08);243-248. www.cdc.gov/mmwr/preview/mmwrhtml/mm6008a4.htm

Prevantics® products are the first and only FDA approved 3.15% Chlorhexidine Gluconate (w/v) and 70% Isopropyl Alcohol (v/v) solution.

The Prevantics® Difference


The efficacy of Prevantics® products closely correlate with dressing change protocols at many facilities.



Clinical Case Study

A leading facility who has implemented Prevantics® products to improve patient outcomes.

The Rush-Copley Medical Center Study resulted in a CLABSI reduction of 59% in year 1 and a 100% reduction by year 4.



*Swabstick and Maxi Swabstick only

³Matocha et al. A Process Improvement Approach to the Elimination of Central Line Associated Bloodstream Infections. Rush-Copley Medical Center, Aurora, IL. October 2011.

⁴Marlowe, Lauren; Mistry, Rakesh D. et al. Blood Culture Contamination Rates after Skin Antisepsis with Chlorhexidine Gluconate versus Povidone-Iodine in a Pediatric Emergency Department. Infection Control and Hospital Epidemiology. February 2010, vol. 31, no. 2.

Prevantics® products are available in a variety of convenient, ready-to-use applicators: Swab, Swabstick and Maxi Swabstick.

PRODUCT BENEFITS

Saves Time	<ul style="list-style-type: none"> • Pre-moistened. No waiting for activation • Unique 2-sided applicators* 	<ul style="list-style-type: none"> • No glass • Easy to open packaging
Saves Money	<ul style="list-style-type: none"> • Effective skin antiseptics will help improve patient outcomes and reduce healthcare costs 	
Improves Outcomes	<ul style="list-style-type: none"> • Up to 7 days of continued antimicrobial activity* • Clinical compendium with summary of independent studies • 14 independent clinical studies demonstrating clinical efficacy with Prevantics® products 	
Meets Evidence-Based Clinical Guidelines	<ul style="list-style-type: none"> • Prevantics® products meet the skin antiseptics guidelines from: <div style="display: flex; justify-content: space-around; align-items: center;">          </div>	

ORDER INFORMATION

Individual Boxes	Reorder No.	Swabs/Swabsticks Per Box	Boxes Per Case	Treatment Area Covered
Prevantics® Swab 1.0 mL Preinjection Indication	B10800	100	10	2.5" x 2.5" in ² (6.25 in ² / 6 x 6 cm)
Prevantics® Swabstick 1.6 mL Preinjection and Preoperative Indications	S40750	50	10	Dry Site: 4" x 4" in ² (16 in ² / 10 x 10 cm) Moist Site: 3" x 5" in ² (15 in ² / 7.5 x 12.5 cm)
Prevantics® Maxi Swabstick 5.1 mL Preinjection and Preoperative Indications	S41950	30	10	Dry Site: 7" x 7" in ² (49 in ² / 18 x 18 cm) Moist Site: 3" x 7.5" in ² (22.5 in ² / 7.5 x 19 cm)

Available for Kits	Reorder No.	Swabs/Swabsticks Per Box	Boxes Per Case	Treatment Area Covered
Prevantics® Swab 1.0 mL Preinjection Indication	B11400	3000	1	2.5" x 2.5" in ² (6.25 in ² / 6 x 6 cm)
Prevantics® Swabstick 1.6 mL Preinjection and Preoperative Indications	S32450	500	1	Dry Site: 4" x 4" in ² (16 in ² / 10 x 10 cm) Moist Site: 3" x 5" in ² (15 in ² / 7.5 x 12.5 cm)
Prevantics® Compact Swabstick 1.6 mL Preinjection and Preoperative Indications	S42850	500	1	Dry Site: 4" x 4" in ² (16 in ² / 10 x 10 cm) Moist Site: 3" x 5" in ² (15 in ² / 7.5 x 12.5 cm)
Prevantics® Maxi Swabstick 5.1 mL Preinjection and Preoperative Indications	S27350	300	1	Dry Site: 7" x 7" in ² (49 in ² / 18 x 18 cm) Moist Site: 3" x 7.5" in ² (22.5 in ² / 7.5 x 19 cm)

*Swabstick and Maxi Swabstick only

PDI
Two Nice-Pak Park
Orangeburg, New York 10962
Customer Service: 800.999.6423
pdihc.com

©2014 Professional Disposables International, Inc. All rights reserved.

