

Prevantics® Swabstick

Skin Antisepsis

3.15% Chlorhexidine Gluconate (w/v) and 70% Isopropyl Alcohol (v/v)
Preinjection & Preoperative Skin Preparation

INSTRUCTIONS FOR USE

- For Item No. S40750: Peel apart the pouch to expose the swabstick.
- For Item No. S42850 (compact swabstick – kit) and Item No. S32450 (swabstick – kit): Grasp the pouch at the notch and tear open straight across.
- Remove the swabstick from package. Do not touch the foam tip.



PEEL APART
Item No. S40750



Item No. S42850
(compact swabstick – kit) **TEAR OPEN**



Item No. S32450
(swabstick – kit)

APPLY

- For dry sites:** Scrub repeatedly back and forth for 15 seconds with each side of the applicator for a total of 30 seconds. Allow to air dry for approximately 30 seconds.
- For moist sites:** Scrub repeatedly back and forth for 1 minute with each side of the applicator for a total of 2 minutes. Allow to air dry for approximately 1.5 minutes.
- Maximum treatment area for one swabstick:
 - Dry Site: 4 x 4 in. (10 x 10 cm.)**
 - Moist Site: 3 x 5 in. (7.5 x 12.5 cm.)**
- Do not blot or wipe dry.
- Discard after single use.



See Drug Facts panel for complete product information.

REORDER # _____ REPLACING ORDER # _____

1-800-999-6423 + Phone: (845) 365-1700 + Fax: (845) 398-5347 + www.PDIHC.com

©2018 PDI.
PDI02189755

PDI
BE THE DIFFERENCE™