

Prevantics® Maxi Swabstick

Skin Antisepsis

3.15% Chlorhexidine Gluconate (w/v) and 70% Isopropyl Alcohol (v/v)
Preinjection & Preoperative Skin Preparation

INSTRUCTIONS FOR USE

- For Item No. S41950: Peel apart the pouch to expose the swabstick.
- For Item No. S27350 (kit): Grasp the pouch at the notch and tear open straight across.
- Remove the swabstick from package. Do not touch the foam tip.



PEEL APART
Item No. S41950



TEAR OPEN
Item No. S27350 (kit)

APPLY

- Prior to surgery or injection, apply swabstick to the procedure site.
- Place one flat side of foam tip to the proposed skin site and prep the skin in vigorous back-and-forth repeated strokes for 1 minute.
- Turn the swabstick over (unused side of the foam tip) and repeat procedure by prepping for 1 minute.
- Allow the prepped area to air dry for 1.5 minutes.
- Maximum treatment area for one swabstick:
 - **Dry Site: 7 x 7 in. (18 x 18 cm.)**
 - **Moist Site: 3 x 7.5 in. (7.5 x 19 cm.)**
- Do not blot or wipe dry.
- Discard after single use.



See Drug Facts panel for complete product information.

REORDER # _____ **REPLACING ORDER #** _____

1-800-999-6423 + Phone: (845) 365-1700 + Fax: (845) 398-5347 + www.PDIHC.com

©2013 PDI.
PDI01136575_DA


PDI
BE THE DIFFERENCE™