

**1 ONE UNIT  
1 ONE DOSE  
1 ONE AND ONLY**



**Unit Dose Non-Aerosol Spray  
20% Benzocaine Oral Anesthetic  
0.5 mL each**

**ORIGINAL WILD CHERRY FLAVOR**

**DIRECTIONS**

<p><b>PRODUCT DIAGRAM</b></p>	<p><b>TEAR</b></p> <p>at notch to open pouch. Remove HurriCaine ONE® from pouch.</p> <p><b>1</b></p>	<p><b>GRASP</b></p> <p>HurriCaine ONE firmly by placing your first and second fingers on the finger grip, while your thumb is touching the base of the plunger.</p> <p><b>2</b></p>	<p><b>POSITION</b></p> <p>patient with mouth open. Aim spray tip toward the back of patient's throat. NOTE: For added support, you may use your other hand or third finger as a fulcrum.</p> <p><b>3</b></p>	<p><b>APPLY</b></p> <p>firm pressure to plunger by pushing thumb forward to initiate spray action. Instruct patient to swallow 1-2 times. Anesthesia is accomplished within 30 seconds and lasts approximately 15 minutes.</p> <p><b>4</b></p>
-------------------------------	--	---	--	--

<b>Drug Facts</b>	
<b>Active ingredient</b> Benzocaine 20% .....	<b>Purpose</b> .....Oral Anesthetic
<b>Uses</b> for the temporary relief of occasional minor irritation and pain associated with sore mouth and throat	
<b>Warnings</b>	
<b>Allergy alert:</b> Do not use if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine, or other "caine" anesthetics.	
<b>When using this product</b> avoid contact with eyes	
<b>Stop use and ask a doctor if</b>	
<ul style="list-style-type: none"> <li>■ sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, swelling, nausea or vomiting</li> <li>■ sore mouth symptoms do not improve in 7 days, or if irritation, pain, or redness persists or worsens</li> </ul>	
<b>Keep out of reach of children.</b> If more than used for pain is accidentally swallowed, get medical help or contact a Poison Control Center immediately.	
<b>Adverse Reactions:</b> Methemoglobinemia has been reported following the use of benzocaine on extremely rare occasions. Intravenous methylene blue is the specific therapy for this condition.	
<b>Directions</b> do not exceed recommended dosage	
<b>age</b>	<b>dose</b>
adults and children 2 years of age and older	use up to 4 times daily or as directed by a dentist or doctor
children under 12 years of age	should be supervised in the use of the product
children under 2 years of age	consult a dentist or doctor
<b>Other information</b> ■ store at 15-30 °C (59-86 °F) ■ do not use if foil pouch has been previously opened	
<b>Inactive ingredients</b> ■ flavor, polyethylene glycol, sodium saccharin	
<b>Questions or comments?</b> 1-800-238-8542 M-F: 8:00 a.m. – 4:30 p.m. ET	



**MAKE THE SWITCH TODAY.**

Contact your preferred dealer or call 1-800-238-8542 to place your order or for more information.  
www.beutlich.com

HurriCaine ONE is a registered trademark of Beutlich® Pharmaceuticals, LLC. HODH 614 1214